



Supplemental Labeling

FOR DISTRIBUTION AND USE ONLY WITHIN IDAHO

EPA Reg. No. 241-337

EPA SLN No.: ID-000003

FOR WEED CONTROL IN PERENNIAL GRASSES GROWN FOR SEED PRODUCTION

This label is valid until December 31, 2015 or until otherwise amended, withdrawn, canceled or suspended.

For application to established (6 or more tillers per plant) Kentucky bluegrass, tall fescue, orchardgrass, perennial ryegrass, bentgrass or fine fescue fields.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered product label for Prowl® 3.3 EC herbicide, EPA Reg. No. 241-337.

This labeling must be in the possession of the user at the time of application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Observe all cautions and limitations on this label and on the labels of products used in combination with **Prowl 3.3 EC**. Do not use **Prowl 3.3 EC** other than in accordance with the instructions set forth on this label. The use of **Prowl 3.3 EC** not consistent with this label may result in injury to crops. Keep containers closed to avoid spills and contamination.

Read the **Precautionary Statement**, **Environmental Hazards**, **Storage and Disposal**, and **Disclaimer** statements appearing on the container label.

GENERAL INFORMATION

Prowl 3.3 EC applied prior to weed emergence will control annual bluegrass, volunteer fescue, and volunteer ryegrass in addition to the weeds listed on the **Prowl 3.3 EC** label. It may be necessary to control weeds that have germinated prior to the application of **Prowl 3.3 EC** with an alternative herbicide that has postemergence activity on those weeds. Weed control can be enhanced by using sequential applications of other registered pesticides.

Prowl 3.3 EC may cause temporary injury to grass stands. Diseases, extremely cold weather, drought, extensive frost heaving, low or high pH, or salinity may weaken strands and make them more susceptible to herbicidal damage.

APPLICATION INSTRUCTIONS

Apply **Prowl 3.3 EC** at the beginning of significant fall rains and prior to weed germination. Uniformly apply 4.8 to 7.2 pints of **Prowl 3.3 EC** per acre in 10 or more gallons of water per acre with ground equipment. The higher application rate should be used where heavy weed

seedling or volunteer seedling infestations are anticipated.

Prowl® 3.3 EC herbicide treatments are most effective when adequate rainfall or overhead irrigation is received within 7 days after application. Effectiveness of **Prowl 3.3 EC** may be increased by removal of excess straw and crop residue by such methods as crew cutting, propane flaming, or open field burning when local regulations allow.

CHEMIGATION APPLICATION

Prowl 3.3 EC may be applied through sprinkler irrigation systems including center pivot, lateral move, side (wheel) roll, solid set, or hand move irrigation systems. For complete instructions and precautions regarding chemigation refer to the main label for **Prowl 3.3 EC**.

Prior to weed germination, uniformly apply 4.8 to 7.2 pints of **Prowl 3.3 EC** per acre through sprinkler irrigation systems.

If soil condition is extremely dry, a pre-chemigation irrigation of $\frac{1}{4}$ - $\frac{1}{2}$ inch is recommended to enhance the incorporation of **Prowl 3.3 EC**. Chemigate as soon as possible after the pre-chemigation irrigation.

RESTRICTIONS AND LIMITATIONS

DO NOT apply more than 7.2 pints of **Prowl 3.3 EC** per crop season.

DO NOT apply if surface water is present.

DO NOT when wind conditions favor drift.

DO NOT harvest treated grass for feed or allow livestock to graze treated areas.

FOLLOW CROP RESTRICTIONS

Any crop labeled for **Prowl 3.3 EC** preplant incorporated (PPI) use can be planted the same year without adverse effect.

For Prowl 3.3 EC use rates up to 5 pints/acre:

Wheat may be planted 12 months following a **Prowl 3.3 EC** application. Other crops may be planted 18 months following an application of **Prowl 3.3 EC**.

For Prowl 3.3 EC use rates up to 7.2 pints/acre:

Crops other than sugar beets, red beets, and spinach may be planted 18 months following application. Sugar beets, red beets, and spinach may be planted 24 months after application.

Refer to the **Prowl 3.3 EC** herbicide label for complete follow crop restrictions.

©2005 BASF Corporation
All rights reserved

Prowl is a registered trademark of BASF.

0000241-00337.20051007.NVA 2005-04-129-0318
Supercedes NVA 2004-04-129-0244

24(c) Registrant:

BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709



The Chemical Company